




INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ



I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.


In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated 28 October 2004





The Patent Office

22OCT03 E846258-3 D01038
P01/7700 0.00-0324551.1

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference

S6729/P709871GB/DGB/KB530

2. Patent application number

(The Patent Office will fill in this part)

0324551.1

21 OCT 2003

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Karo Bio AB
Novum
SE-141 57 Huddinge
Sweden

Patents ADP number (if you know it)

8214421001

If the applicant is a corporate body, give the country/state of its incorporation

SE

4. Title of the invention

NOVEL COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

DAVID GARDNER BANNERMAN
WITHERS & ROGERS
Goldings House
2 Hays Lane
London
SE1 2HW

Patents ADP number (if you know it)

1776001 /

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

YES

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form

Description 56
Claim(s) 13
Abstract 1
Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents
Translations of priority documents
Statement of inventorship and
right to grant of a patent (Patents Form 1/77)
Request for preliminary examination
and search (Patents Form 9/77) 1
Request for substantive examination
(Patents Form 10/77)
Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature *W. G. Bannerman*

Date 21 October 2003

12. Name and daytime telephone number of
person to contact in the United Kingdom

David G Bannerman

01926 336111

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500 505.
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- e) Once you have filled in the form you must remember to sign and date it.
- f) For details of the fee and ways to pay please contact the Patent Office.



5

NOVEL COMPOUNDS

Field of Invention

10 This invention relates to novel compounds which are antagonists or partial antagonists of the androgen receptor and the use of such compounds for therapeutic purposes.

Background of Invention

15

The androgen receptor (AR) is a member of the steroid hormone nuclear receptor family of ligand activated transcription factors. This group includes estrogen, progesterone, mineralocorticoid, and glucocorticoid receptors all of which are activated by endogenous steroid hormones to control the expression of responsive genes. The hormone receptors share a modular structure consisting of a variable amino-terminal domain (NTD), a highly conserved DNA-binding domain (DBD), and a carboxy-terminal ligand-binding domain (LBD). The DNA-binding domain generates much of the transcriptional specificity due to its ability to discern different DNA response elements with the promoter regions of target genes. The LBD is required for ligand dependent transcriptional activity containing both the hormone-binding pocket and an important transcriptional activation functional region (AF2) required for recruitment of coactivators and the cellular transcriptional machinery.

20

25

Regulation of nuclear receptor activity resides predominantly in the binding of the hormone ligand within the LBD. The amino acids lining the interior of the hormone-binding cavity define the selectivity of the receptor for its hormone. This allows AR to discriminate between the natural ligands and non-natural ligands.

30

5 The amino acids and the "space" they define as the hormone-binding cavity can be exploited in synthesizing modulators that are highly receptor selective. These interactions between the endogenous hormone and amino acid residues within the ligand-binding cavity induce conformational changes that are distributed throughout the entire receptor structure. It is these conformational changes that lead to the dissociation of chaperone
10 proteins that stabilize the receptors in the absence of ligand and the association of coactivator proteins. A liganded receptor devoid of its chaperone proteins is able to dimerize, translocate, recruit coactivators, and initiate transcription.

 The natural ligand for the androgen receptor, androgen, is produced in both men and
15 women by the gonads, adrenal glands and locally in target tissues. The levels of androgens secreted by the gonads are tightly regulated by a feedback mechanism involving the hypothalamus and pituitary.

 In men, androgens are necessary for masculinization and fertility. However,
20 systemic androgen excess causes testicular atrophy and infertility. Androgens may also contribute to lipid abnormalities, cardiovascular disease and psychological abnormalities. Local androgen excess is implicated in the pathogenesis of male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne. The physiologic role of androgens in women is not well understood, but these steroids do play a role in the
25 development of normal body hair and libido. In women, relative androgen excess causes hirsutism (excessive hair growth), amenorrhea (abnormal loss or suppression of menses), acne and male pattern baldness.


 The risk of developing prostate cancer increases dramatically with age. More than
30 75% of prostate cancer diagnoses are in men over the age of 65, and the prevalence of clinically undetectable prostate cancer in men over 80 years old is as high as 80%. It remains unclear as to the exact cause of prostate cancer, however, it is widely accepted that androgens can increase the severity and the rate of progression of the disease.

- 5 Androgen deprivation therapy has been the basis for prostate cancer therapy since 1941 when castration was shown to have beneficial effects on advanced stages of the disease. Hormonal intervention is currently based on disrupting the hypothalamus-pituitary-gonadal feedback mechanism to control the levels of endogenous androgens from the testes. Antandrogens are incorporated in later stage therapies to work at the level of the
- 10 androgen receptor itself, blocking residual androgens from adrenal sources. In spite of these treatments, there exists a need for an improved therapy of diseases linked to disturbances in the activity of the androgen receptor.

SUMMARY OF THE INVENTION

15

The present invention provides use as a medicament of a compound according to Formula I



Formula I

20

in which;

- R₁ and R₂ are the same or different and independently selected from the group consisting of; hydrogen, halogen, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkoxy, C₁-C₁₀ alkenoxy, C₁-C₁₀ alkenoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ alkenylthio,
- 25 C₁-C₁₀ alkynylthio, C₆-C₁₀ arylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₆-C₁₀ aryl, or C₃-

- 5 C₂₀ heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different;

- R₃ and R₄ are the same or different and independently selected from the group consisting of; hydrogen, halogen, keto, C₁-C₂₀ alkyl, C₃-C₇ cycloalkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ alkenoxy, C₁-C₄ alkynoxy, C₁-C₄ alkylthio, C₁-C₄ alkenylthio, C₁-C₄ alkynylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₆-C₁₅ aryl, or C₅-C₂₀ heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different;

R₅ is chosen from the group consisting of; nitro, cyano, acetic acid, halogen, sulphonic acid, aldehyde, carboxylic acid or ester, phosphonic acid, or ester;

- 20 R₆ is chosen from the group consisting of; C₁-C₅ alkyl, halogen, CN, CO₂H, CHF₂, CH₂F, or CF₃;

R₈ is chosen from the group consisting of; H, C₁-C₅ alkyl, halogen, CHF₂, CH₂F or CF₃;

- 25 X is chosen from the group consisting of; -NH-, -O-, -S-, -SO-, -SO₂-, -Se-, -Te- or -S-S-

Y is chosen from the group consisting of; OH, NH₂, unbranched, branched or cyclic C₁-C₅ alkyl, unbranched, branched or cyclic -NH(C₁-C₈); unbranched, branched or cyclic N(C₁-C₈)₂, -NH(C₆ aryl), -N(C₆ aryl)₂, -NH(C₁-C₅ heteroaryl), and -N(C₁-C₅ heteroaryl)₂ or a bioisosteric equivalent;

Z is chosen from the group consisting of; CR₇, N, or O;

5 R_7 is H, or C_1-C_5 alkyl;

R^a represents a member selected from the group consisting of; hydrogen, halogen, -CN, CO_2H , CHO, NO_2 , $-NH_2$, $-NH(C_1-C_4)$; $N(C_1-C_4)_2$, $-NH(C_6\text{ aryl})$, $-N(C_6\text{ aryl})_2$, $-NH(C_1-C_5\text{ heteroaryl})$, and $-N(C_1-C_5\text{ heteroaryl})_2$ or a bioisosteric equivalent; or a

10 pharmaceutically acceptable salt thereof;

A preferred compound is according to formula I, wherein R_1 or/and R_2 are H, (S)-methyl, methyl, (R)-ethyl, (S)-ethyl, ethyl, (R)-propyl, (S)-propyl, propyl, (S)-butyl, (S)-1-methyl-propyl, (S)-2-methyl-propyl, (R)-isopropyl, (S)-isopropyl, isopropyl,
15 cyclopentyl, $-(CH_2)_2SMe$, (R)- CH_2SCH_2Ph , (S) - benzyl, 4-chloro-benzyl, (S)-methyl-1-H-indole or (S)-phenyl;

Further preferred is a compound according to formula I, wherein R_3 is chosen from the group consisting of; hydrogen, methyl, ethyl, phenyl, 3-hydroxy phenyl, 4-hydroxy
20 phenyl, or keto.

Further preferred is a compound according to formula I, wherein R_4 is H, methyl, or keto.

Further preferred is a compound according to formula I, wherein R_5 is NO_2 , CN, CH_2CN
25 or CO_2H ;

Further preferred is a compound according to formula I, wherein R_6 is Me, or CF_3 ;

Further preferred is a compound according to formula I, wherein R_7 is H or Me;
30

Further preferred is a compound according to formula I, wherein R_8 is H or methyl;

Further preferred is a compound according to formula I, wherein X is NH;

5

Further preferred is a compound according to formula I, wherein Y is H, -OH, -OMe, -N(CH₂CH₃)₂, piperidine, or 4-nitro-2-ylamino;


Further preferred is a compound according to formula I, wherein Z is CR₇ or N;

10









Even more preferred is a compound according to formula I, chosen from the group consisting of;










- 2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol;
- 15 [1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol;
- (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol;
- (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol;
- 2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol;
- [1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol;
- 20 (S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol;
- [1-(6-Methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol;
- (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-2-phenyl-ethanol;
- (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol;
- (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol;
- 25 (DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
- (S)-2-(6-Methyl-5-nitro-2-pyridin-ylamino)-propionic acid;
- (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
- 2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol;
- (S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol;
- 30 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile;
- 4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
- (S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
- (R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;

- 5 (S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
 [4-((S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 [4-((R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 [4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile;
 10 6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile;
 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile;
 and the compounds showed in the following table (The substituents, R₉, R₆, and Z, are
 shown in the table, and are all substituents in the following formula II. In formula II, the
 NO₂ group corresponds to the substituent R₅ in formula I, and R₉ is composed of the
 15 moieties X R₁ R₂ Y R₃ R₄ of Formula I as defined above, where X is -NH-






















Formula II

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH

R9	R6	Z		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	CH

CC(C)[C@H](N)CO CH_3

CH

 CH_3

CH

 CH_3

CH



CH

CE



CE

CE



CE

C



C

C



C

1 c



C

1.



C

1.



1

1

- 5 Since the compounds are shown to be mainly antagonists for the androgen receptor, a preferred use is the use of the composition above for treating a disease which is caused by an increase in androgen receptor activity.

10 Even more preferred is the use of the composition above for treating a disease which is chosen from the group consisting of; prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.


15 The present invention also provides the use of a compound according to the above for manufacturing a medicament to be used in the treatment of a disease caused by a disturbance in the activity of the androgen receptor.

- 20 Since the compounds according to the invention are mainly antagonists, these are mainly suited to treat diseases that are caused by an increase in androgen receptor activity.

25 A specific disease that would be amenable for treatment by the present invention is a disease chosen from the group consisting of; prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.

- 30 The compounds of the present invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more other agent(s) active in the therapeutic areas described herein.

- 5 According to another aspect of the invention there is provided a compound as defined in Formula I, provided that the compound is not the compound according to the formula;



- 10 The specific compound above is known in the prior art as an intermediate compound in the manufacture of other compounds.

DETAILED DESCRIPTION OF THE INVENTION

- 15 The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

- 20 The term "androgen receptor ligand" as used herein is intended to cover any moiety, which binds to an androgen receptor. The ligand may act as an antagonist, or as a partial antagonist.

A compound being a "partial antagonist" is a compound with both agonistic and antagonistic activity.

- 25 The term "alkyl" as employed herein alone or as part of another group refers to an acyclic straight or branched chain radical, containing 1 to about 10 carbons, preferably 1 to 6 carbons in the normal chain, i.e. methyl, ethyl, propyl, isopropyl, sec-butyl, tert-butyl, pentyl, hexyl, octyl. When substituted alkyl is present, this refers to an unbranched or branched alkyl group, which groups may be the same or different at any available
30 point, as defined with respect to each variable.

5 The term "alkyl" includes an alkyl group optionally substituted with one or more functional groups which are commonly attached to such chains, such as, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, hydroxy, cyano, nitro, amino, halo, carboxyl or alkyl ester thereof and/or carboxamide.

10 The term "alkenyl" as employed herein alone or as part of another group refers to a straight or branched chain radical, containing 2 to about 10 carbons, preferably 2 to 6 carbons i.e. ethenyl, propenyl, butenyl, allyl.

 The term "allyl" refers to $\text{H}_2\text{C}=\text{CH}-\text{CH}_2$.

15 The term "alkynyl" as employed herein alone or as part of another group refers to a straight or branched chain radical, containing 2 to about 10 carbons, preferably 2 to 6 carbons i.e. ethynyl, propynyl, butynyl, allyl.

20 The term "aryl" as employed herein alone or as part of another group refers to substituted and unsubstituted aromatic ring system. The term aryl includes monocyclic aromatic rings, polycyclic aromatic ring system and polyaromatic ring systems. The polycyclic aromatic and polyaromatic ring systems may contain from two to four, more preferably two to three rings. Preferred aryl groups include 5- or 6- membered ring
25 systems.

 The term "heteroaryl" refers to optionally substituted aromatic ring system having one or more heteroatoms such as, for example, oxygen, nitrogen and sulfur. The terms heteroaryl includes five- or six-membered heterocyclic rings, polycyclic heteroaromatic
30 ring system and polyheteroaromatic ring systems. The poly heterocyclic aromatic and poly heteroaromatic ring systems may contain from two to four, more preferably two to three rings. The term hetero aryl includes ring system such as pyridine, quinoline, furan, thiophene, pyrrole, imidazole and pyrazole.

5

The term "alkoxy" as employed herein alone or as part of another group refers to an alkyl ether wherein the term alkyl is as defined above. Examples of alkoxy radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy and the like.

- 10 The term "aryloxy" as employed herein alone or as part of another group refers to an aryl alkyl ether, wherein the term aryl is as defined above. Examples of aryloxy radicals include phenoxy, benzyloxy and the like.

- 15 The term "alkylthio" as employed herein alone or as part of another group refers to an alkyl thio wherein the term alkyl is as defined above. Examples of alkylthio radicals include methane thiol, ethane thiol, propane thiol and the like.

- 20 The term "alkylsulphone" as employed herein alone or as part of another group refers to an alkylsulphone wherein the term alkyl is as defined above. Examples of alkylsulphone radicals include methanesulphone, ethanesulphone, propanesulphone and the like.

- 25 The term "alkylsulphoxide" as employed herein alone or as part of another group refers to an alkylsulphoxide wherein the term alkyl is as defined above. Examples of alkylsulphoxide radicals include methanesulphoxide, ethanesulphoxide, propanesulphoxide and the like.

- 30 The term "cycloalkyl" as employed herein alone or as part of another group refers to saturated cyclic hydrocarbon groups or partially unsaturated cyclic hydrocarbon groups, independently containing one carbon-to-carbon double bond. The cyclic hydrocarbon contains 3 to 4 carbons. It should also be understood that the present invention also involve cycloalkyl rings where 1 to 2 carbons in the ring are replaced by either -O-, -S- or -N-, thus forming a saturated or partially saturated heterocycle. Examples of such rings

- 5 are aziridine, thiiranes and the like. Preferred heterocyclic rings are 3-membered, which may be optionally substituted by 1, 2 or 3 groups of R^a which groups may be the same or different through available carbons as in the case of "alkyl". Preferred cycloalkyl groups include 3 carbons, such as cyclopropyl, which may be optionally substituted by 1, 2 or 3 groups of R^a which groups may be the same or different through available carbons as in the case of "alkyl".
- 10

The term "halogen" refers to fluorine, chlorine, bromine and iodine. Also included are carbon substituted halogens such as $-CF_3$, $-CHF_2$, and $-CH_2F$

- 15 The compounds of the present invention can be present as salts, which are also within the scope of this invention. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred. If the compounds of the invention have, for example, at least one basic center, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted, for example, by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, (for example aspartic or glutamic acid or lysine or arginine), or benzoic acid, or with organic sulfonic acids, such as (C_1-C_4) alkyl or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methyl- or p-toluene- sulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center.
- 20
- 25
- 30 The compounds of the invention having at least one acid group (e.g. $COOH$) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine,

5 piperidine, pyrrolidine, a mono, di or tri-lower alkylamine, for example ethyl, tertbutyl, diethyl, diisopropyl, triethyl, tributyl or dimethyl-propylamine, or a mono, di or trihydroxy lower alkylamine, for example mono, di or triethanolamine. Corresponding internal salts may furthermore be formed. Salts that are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free
10 compounds of the invention or their pharmaceutically acceptable salts, are also included. Preferred salts of the compounds of the present invention which contain a basic group include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate or nitrate. Preferred salts of the compounds of formula I which contain an acid group include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

15

a27
The compounds according to the invention may also have prodrug forms. Any compound that will be converted in vivo to provide the bioactive agent (i.e., the compound of formula I) is a prodrug within the scope and spirit of the invention. Such prodrugs are well known in the art and a comprehensive description of these may be
20 found in: (i) *The Practice of Medicinal Chemistry*, Camille G. Wermuth et al., Ch 31, (Academic Press, 1996); (ii) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985); and (iii) *A Textbook of Drug Design and Development*, P. Krogsgaard-Larson and H. Bundgaard, eds. Ch 5, pgs 113 – 191 (Harwood Academic Publishers, 1991).

25

Embodiments of prodrugs suitable for use in the present invention include lower alkyl esters, such as ethyl ester, or acyloxyalkyl esters such as pivaloyloxymethyl (POM).

30

The compounds according to the present invention are preferably administered in a therapeutically effective amount. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent to treat or prevent a condition treatable by administration of a composition of the invention. That amount is the amount sufficient to exhibit a detectable therapeutic or preventative or ameliorative effect. The effect may include, for example, treatment or prevention of the conditions listed herein.


- 5 The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition being treated, recommendations of the treating physician, and the therapeutics or combination of therapeutics selected for administration.

10 Scheme 1-6 outlines the synthetic routes used for preparing the compound according to Formula I

Scheme 1

- 15 Synthetic routes to these compounds can be visualized by the skilled person and the present synthetic route is not limiting for the invention. 4-Fluoro-1-nitro-2-trifluoromethyl-benzene (1a) and 4-fluoro-2-methyl-1-nitro-benzene (1b) were employed as starting material in scheme-1 and is commercially obtainable.

- 20 Scheme 1 depicts a synthesis of compounds of formula I in which R_6 is CF_3 and Me and is connected to phenyl ring. Condensation of compound (1a) with different β -amino alcohols and di-isopropyl ethylamine in DMSO gave compound 3 (examples 1-4) in quantitative yield. The reactions were performed in a microwave oven at elevated temperature for a short time. Compound (1b) was used for producing the compound 3 (examples 5-7) and similar conditions were adopted as in examples 1-4. An alternative method was used for the preparation of example-5. The reaction according to the
- 25 alternative method was performed by heating the compound (1b) and β -amino alcohol in pentanol in a sealed tube.




Scheme 1

Scheme 2

Compounds 9 (examples 8-15) were prepared from starting material 6-chloro-3-nitro-2-picoline (compound 4). Starting material was synthesized in three steps starting with compound 6-amino-2-picoline using the literature procedure. Nitration of 6-amino-2-picoline was accomplished by concentrated sulphuric acid (H_2SO_4) and concentrated nitric acid (HNO_3) and provided 6-amino-3-nitro-2-picoline (Baumgarten, H. E. and Chien Fan Su, H. *JACS* 74 (1952) 3828; Parker, E. D. and Shive, W. *JACS* 69 (1947) 63). Treatment of 6-amino-3-nitro-2-picoline with sodium nitrite provided 6-hydroxy-3-nitro-2-picoline, which, when reacted with PCl_5 and $POCl_3$, provided 6-chloro-3-nitro-2-picoline (Baumgarten, H. E. and Chien Fan Su, H. *JACS* 74 (1952) 3828).


Scheme 2 shows the synthesis of compounds of formula I in which Z is N and R_1 is H. Condensation of 6-Chloro-3-nitro-2-picoline and 2-amino-2-methyl-propan-1-ol in 1-pentanol and the mixture refluxed under inert atmosphere gave compound 9 (example-8) as yellow crystals. 6-Chloro-3-nitro-2-picoline can also be purchased as commercial starting material. The reaction time was reduced by using a microwave oven. Condensation of compound 7 with different β -amino alcohols (8) in the microwave provided compound 9 (examples 9-13) in quantitative yield. Synthetic routes to these compounds can be visualized by the skilled person. Reaction of compound (10) with L-alanine provided compound 11 (example-14). Reduction of the acid compound (11) by a reducing agent such as lithium aluminum hydride (LAH) produced compound 9 (example 15).



Scheme 2

Scheme 3

- 10 Synthesis of compounds according to formula I, in which R_6 and R_7 are Me and connected to the phenyl ring is shown in Scheme-3. 4-Fluoro-2, 3-di-methyl-1-nitro-benzene (13) was employed as starting material in scheme-3, which was produced by the nitration of compound (12) with fuming nitric acid in acetic anhydride in one step. Condensation of 2, 3-dimethyl-fluoro-benzene with β -amino alcohols at higher temperature gave compound 14 (example 16).
- 15



5

12


13

14

Scheme 3

Scheme 4

Scheme 4 depicts a synthesis of compounds of formula I in which R₆ and R₈ are Me and connected to the phenyl ring. Condensation of compound (15) with (S)-2-amino-
 10 butan-1-ol and di-isopropyl ethylamine in DMSO gave compound 16 (examples 17). The reaction was performed in a microwave oven.



15

16


Scheme 4

15

Scheme 5

Reduction of nitro compound to amine was accomplished by the treatment of sodium
 thiosulphate with ethanol. After work-up the amines were used for the next step without
 20 any further purification. Reaction of amine with potassium cyanide and copper cyanide in
 water gave compound 19 (examples 26-28). (Clive, D. L. *et al. JOC* 52 (1987) 1339-42
 and Vogel expt. 6.76). Some other examples of compound 19 were made by an
 alternative method utilizing a microwave oven. Similar reaction conditions as those used
 in scheme-1 and scheme-2 provided compound 19 (examples 18-22).
 25 Conversion of the nitrile form of compound 19 to benzoic acid compound 20 (example
 87) was performed in a refluxed aqueous sodium hydroxide solution in methanol.

5




Scheme 5

10 Scheme 6

Scheme 6 depicts a synthesis of compounds of formula I in which R_3 and R_4 are Me and is connected to the alkyl chain. Condensation of 6-chloro-3-nitro-2-picoline with glycine methyl ester hydrochloride and triethyl amine in DMSO gave compound 22 (example 88). Compound 22 was treated with methyl magnesium bromide and after HPLC purification gave compound 23 (example 89).

15




Scheme 6

5

EXAMPLES

The following Examples represent preferred embodiments of the present invention. However, they should not be construed as limiting the invention in any way. The ¹H NMR spectra were consistent with the assigned structures. Mass spectra were recorded on a Perkin-Elmer, API 150Ex spectrometer, with turbo "ion spray" on negative ion mode (ES-1) or positive (ES+1), using a Zorbax SB-C8 column (LC-MS). The microwave reactions were performed in a Personal Chemistry Emrys Optimizer.

15

Example 1

20


2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (1.226 g, 5.86 mmol) was dissolved in 7 mL DMSO and 2-amino-2-methyl-propan-1-ol (784 mg, 8.795 mmol) was added, followed by diisopropyl ethylamine (DIPEA) (985 mg, 7.622 mmol). The reaction was heated to 180 °C for 900 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed three times with an aqueous solution of ammonium chloride (NH₄Cl). The organic phase was collected, dried with MgSO₄ (anhydrous) and filtered. The dry organic phase was evaporated *in vacuo*. The crude product was a bright yellow powder. The crude product was purified on a silica column with 5:1 n-heptane: EtOAc as mobile

30

5 phase. This gave 1.1 g (68 %) of 2-methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol as a yellow solid. $M/Z = 278$

Example 2




10

[1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (122 mg, 0.583 mmol) was coupled with (1-
15 amino-cyclopentyl)-methanol (101 mg, 0.875 mmol), DIPEA (90.5 mg, 0.700 mmol) in
DMSO 0.8 mL, using the same procedure as described in Example-1. This gave 120.5 mg
(68%) of [1-(4-nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol as a yellow
powder. $M/Z = 304$.

Example 3

20




(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol

25 4-Fluoro-1-nitro-2-trifluoromethyl-benzene (119 mg, 0.569 mmol) was coupled with (S)-
2-amino-3-phenyl-propan-1-ol (129 mg, 0.854 mmol), DIPEA (88 mg, 0.683 mmol) in
DMSO 0.8 mL using the same procedure as described in Example-1. This gave 112 mg

- 5 (58%) of (S)-2-(4-nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol as yellow crystals. $M/Z = 340$.

Example 4




10

(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol

- 4-Fluoro-1-nitro-2-trifluoromethyl-benzene (122 mg, 0.583 mmol) was coupled with (S)-2-amino-butan-1-ol (78 mg, 0.875 mmol), DIPEA (91 mg, 0.700 mmol) in DMSO 0.8 mL using the same procedure as described in Example-1. This gave 107 mg (67%) of (S)-2-(4-nitro-3-trifluoromethyl-phenylamino)-butan-1-ol as yellow oily crystals. $M/Z = 278$.

Example 5



20

2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol


- Method-A: 4-Fluoro-2-methyl-1-nitro-benzene (113 mg, 0.728 mmol) was coupled with 2-amino-2-methyl-propan-1-ol (84 mg, 0.947 mmol), DIPEA (122 mg, 0.947 mmol) in DMSO 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 72 mg (44 %) of 2-methyl-2-(3-methyl-4-nitro-phenylamino)-propan-1-ol as yellow powder. $M/Z = 224$.

30

- 5 Method-B: 4-Fluoro-2-methyl-1-nitro-benzene (2.33 g, 15 mmol) and 2-amino-2-methylpropanol (2.67 g, 30 mmol) were heated with stirring at 160°C in a sealed tube overnight. The reaction mixture was diluted with EtOAc and purified by flash chromatography (dry application; 14% EtOAc in hexane → EtOAc) to afford 2.85 g (85%) of the 2-methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol.

10

Example 6




15

[1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol

- 4-Fluoro-2-methyl-1-nitro-benzene (107 mg, 0.689 mmol) was coupled with (1-amino-cyclopentyl)-methanol (103 mg, 0.897 mmol), DIPEA (116 mg, 0.897 mmol) in DMSO 20 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 nheptane: EtOAc as mobile phase. This gave 76 mg (44 %) of [1-(3-methyl-4-nitro-phenylamino)-cyclopentyl]-methanol as a yellow powder. $M/Z = 250$.

25

Example 7




(S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol

30

- 5 4-Fluoro-2-methyl-1-nitro-benzene (102 mg, 0.658 mmol) was coupled with (S)-2-amino-butan-1-ol (76 mg, 0.855 mmol), DIPEA (111 mg, 0.855 mmol) in DMSO 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 85 mg (58 %) of (S)-2-(3-methyl-4-nitro-phenylamino)-butan-1-ol as yellow oil. $M/Z = 224$.

10

Example 8

- 15 2-Methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol


- (a) Conc. H_2SO_4 (140 ml) was cooled in an ice-salt bath and molten 6-amino-2-picoline (30 g, 0.277 mol) was added in portions with good stirring. To this brown, viscous solution which was maintained at $0^\circ C$ was added a cooled ($0^\circ C$) mixture of conc. H_2SO_4 (21 ml) and conc. HNO_3 (21 ml) drop wise over a period of approx. 1.5 hrs. The red-orange reaction mixture was stirred for an additional hour at $0^\circ C$ and then allowed to warm slowly to room temperature over night. The brown solution was heated at $60^\circ C$ (oil bath) for 1 hr followed by 1hr at $100^\circ C$ (carefully controlled temperature). The reaction mixture was cooled to $0^\circ C$ (ice bath), poured over cracked ice and neutralised by addition of a concentrated aqueous NaOH solution. The yellow precipitate was filtered and washed well with ice-water. (The filtrate was put in the refrigerator; additional product was precipitated together with the salts.) The yellow product was suspended in water and divided into two portions, each of them subjected to steam distillation in turn. The yellow reaction mixture became more "transparent" after some hrs, but the collected steam, containing 4-amino-3-nitro-2-picoline, was still yellow after 6 hrs. The steam distillation was stopped after 8 hrs, the residual part of the reaction mixture was filtered and

30

- 5 evaporated to dryness. ¹HNMR (D₂O) showed a mixture of 2-3 compounds. The mixture was washed with; CHCl₃, EtOH (x 2) and CHCl₃ leaving 20.4 g (48%) of pure 6-amino-3-nitro-2-picoline.
- (b) 6-Amino-3-nitro-2-picoline (20 g, 0.131 mol) was suspended in a mixture of conc. H₂SO₄ (23.7 ml) and water (335 ml). More conc. H₂SO₄ (20 ml) was added under ice-cooling, but the amine did not dissolve completely. The suspension was added in ice (100 g) before a solution of NaNO₂ (13.53 g, 0.196 mol) in water (40 ml) was added drop wise. Gas evolution was observed. The brown suspension was stirred at 10°C for 1 hr, filtered and washed with water. The brown product was dried (freeze dryer) to achieve 15.78 g (78 %) of 6-hydroxy-3-nitro-2-picoline.
- 15 (c) To 6-Hydroxy-3-nitro-2-picoline (15.73 g, 0.102 mol) was added PCl₅ (5.73 g, 0.027 mol) and POCl₃ (2.9 ml, 0.032 mol). This mixture was heated at 110-115°C for 3 hrs. However, the amount of POCl₃ added was only enough to moisten the starting material. More POCl₃ (3 ml) was added, the reaction mixture heated at 110-115°C but only
- 20 sublimation of PCl₅ (100°C) was observed. DMF (5 ml) was added and the solution was heated at 115°C for 5 hrs, cooled and poured into a slush of ice and water. A beige product precipitated and the water suspension was stirred for 48 hrs. The brown precipitate was filtered off and washed with water. Purification by dry-flash dichloromethane yielded 10.93 g (62 %) of 6-chloro-3-nitro-2-picoline.
- 25 (d) 6-Chloro-3-nitro-2-picoline (6.055 g, 35.1 mmol) and 2-amino-2-methyl-propan-1-ol (6.2 g, 73.7 mmol) were suspended in 1-pentanol (30 ml) and the mixture refluxed under inert atmosphere overnight. The thin layer chromatography (dichloromethane 4/EtOAc 1) revealed some remaining starting material, so the reaction was refluxed for another 3.5
- 30 hrs. The reaction mixture was cooled and water was added under stirring. A sticky, yellow precipitate was filtered off, washed well with water and dried. The crude product (6.04 g) was re-crystallised from either pentane-acetone or dichloromethane. Collecting

- 5 the crops furnished 5.71 (72 %) of 2-methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol as yellow crystals. $M/Z = 225$.

Example 9




10

[1-(6-Methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol

- 6-Chloro-3-nitro-2-picoline (22 mg, 0.13 mmol) was coupled with (1-amino-
 15 cyclopentyl)-methanol (31 mg, 0.27 mmol), triethylamine (0.025 mL, 0.18 mmol) in 2-pentanol (1 mL). The reaction was heated to 180 °C for 2 h in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed with NaHCO_3 . The organic phase was collected, dried with anhydrous MgSO_4 and filtered. The dry organic phase
 20 was evaporated and purified on a silica column with 5:1 n-Heptane: EtOAc as mobile phase. This gave 9 mg (28%) of [1-(6-methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol as a yellow solid. $M/Z = 251$

Example 10



(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino) 2-phenyl-ethanol


30

- 6-Chloro-3-nitro-2-picoline (22 mg, 0.13 mmol) was coupled with (2-amino-2-phenyl)-

- 5 propanol (34 mg, 0.25 mmol) in triethylamine (0.030 mL, 0.25 mmol) in DMSO (1 mL). The reaction was heated to 140 °C for 1200 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed with NH₄Cl (aq) three times. The organic phase was collected, dried with anhydrous MgSO₄ and filtered. The dry organic phase was
- 10 evaporated and purification on silica column with 5:1 n-Heptane: EtOAc gave 22 mg (63%) of (R)-2-(6-methyl-5-nitro-pyridin-2-ylamino) 2-phenyl-ethanol as a yellow solid. M/Z = 273.

Example 11

15




(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol.

20

- 6-Chloro-3-nitro-2-picoline (30 mg, 0.17 mmol) was coupled with (S)-2-amino-3-phenyl-propan-1-ol (32 mg, 0.21 mmol), sodium acetate (28 mg, 0.34 mmol) in EtOH (2 mL). The reaction was heated in a microwave oven for 20 min at 130 °C and then additionally 20 minutes at 150 °C. The reaction was quenched with a saturated aqueous solution of
- 25 NaHCO₃ and extracted with EtOAc and evaporated. Purification on a silica column with a gradient solution of heptane: EtOAc gave 24 mg (48%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol as a yellow solid. M/Z = 287.

5


Example 12

(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol

10

6-Chloro-3-nitro-2-picoline (30 mg, 0.17 mmol) was coupled with (S)-2-amino-butan-1-ol (32 mg, 0.21 mmol), and sodium acetate (28 mg, 0.34 mmol) in EtOH (2 mL) using the same procedure as described in Example-13. This gave 21 mg (53%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol as a yellow solid. $M/Z = 225$.

15


Example 13

(DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

6-Chloro-3-nitro-2-picoline (50 mg, 0.29 mmol) was coupled with (DL)-2-amino-3-(4-chloro-phenyl)-propan-1-ol (103 mg, 0.55 mmol), in triethylamine (0.077 mL, 0.55 mmol) in DMSO (1 mL) using the same procedure as described in Example-1 but at 140 °C. This gave 23 mg (45%) of (DL)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-3-(4-chloro-phenyl)-propan-1-ol as a yellow solid. $M/Z = 321$.

5

Example 14



(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid


10

6-Chloro-3-nitro-2-picoline (62 mg, 0.36 mmol) was coupled with L-alanine (80 mg, 0.90 mmol) and sodium acetate (78 mg, 0.95 mmol) in DMSO 1 mL. The reaction was heated to 140 °C for 600 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The crude mixture was treated with a saturated aqueous solution of NH_4Cl . The reaction mixture was acidified to pH 4 (HCl, 1M). The crude reaction mixture was extracted with EtOAc, and the combined organic layers were washed with water and brine. Purification on silica using a mobile phase CH_2Cl_2 -MeOH-HOAc gave 60 mg (74%) of (S)-2-(6-methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid as a yellow solid. $M/Z = 225$.

15

20

Example 15




25 (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid (60 mg, 0.27 mmol) was added to a nitrogen-purged flask with LiAlH_4 (27 mg, 0.71 mmol). The reaction mixture was refluxed for 2 h and then allowed to reach room temperature and then quenched by

- 5 sequentially adding H₂O (1 mL), NaOH (1M, 1 mL) and H₂O (1 mL). The slurry was centrifuged and the precipitated aluminum salts were washed with dichloromethane. The combined filtrates were evaporated and purification of the residue on a silica column with heptane- EtOAc (3:2) gave 13 mg (22%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol as a yellow solid. M/Z = 211.

10

Example 16




- 15 2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol

- Fuming nitric acid (1.4 g, 20.3 mmol) was cooled to 0°C and acetic anhydride (2.89 g, 28.4 mmol) was added. This solution was added to a cold (0°C) solution of 3-fluoro-1,2-dimethylbenzene (1.0 g, 8.1 mmol) in acetic anhydride (4 ml) over 10 min. The reaction mixture was stirred for 25 min, poured slowly over ice and the water solution extracted with EtOAc (x 3). The collected organic phase was washed with diluted saturated aqueous solution of NaHCO₃ followed by brine before evaporation to dryness. The residue was flash purified on a silica gel column using hexane as a mobile phase to give 2,3-dimethyl-4-fluoro-1-nitro-benzene 0.74 g (54%) as a yellow oil which crystallised upon standing.

- 25 The fluoride (0.576 g, 3.4 mmol) was mixed with 2-amino-2-methylpropanol (0.61 g, 6.8 mmol) in a tube, and the tube was sealed before immersing it into an oil bath and heating at 160°C for 5 days. TLC (Hexane) showed remaining starting material. The reaction mixture was cooled and diluted with EtOAc before purification by flash silica gel chromatography (dry application; 6:4 hexane and EtOAc) to give 0.34 g (59% recovery) of the starting material 2,3-dimethyl-4-fluoro-1-nitro-benzene and 0.20 g (61% based on

- 5 recovered starting material) of the 2-(2,3-dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol. $M/Z = 238$.

Example 17




10

(S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol

- (S)-2-Amino-butan-1-ol (41 mg, 0.461 mmol) was dissolved in DMSO (800 μ L) and
 15 DIPEA (80 μ L, 0.461 mmol) added. 4-Fluoro-2-trifluoromethyl-benzonitrile (60mg,
 0.354 mmol) was added and the reaction mixture was heated to 160 $^{\circ}$ C for 900 seconds in
 a microwave oven (Parameters: High absorbance, Fixed Holding time, pre-stirring 25
 sec). The reaction mixture was then diluted with EtOAc and washed with an aqueous
 solution of NH_4Cl . The organic phase was then dried and evaporated *in vacuo*. The crude
 20 product was purified on silica column with 3:1 n-heptane:EtOAc as the mobile phase.
 This provided 22 mg (26 %) of (S)-2-(3,5-dimethyl-4-nitro-phenylamino)-butan-1-ol.
 $M/Z = 238$

Example 18

25




4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile

- 5 2-Amino-2-methyl-propan-1-ol (25 mg, 0.275 mmol) was dissolved in 0.7 mL DMSO and DIPEA (36 mg, 0.275 mmol) was added. 4-fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was then added and the reaction was heated to 140 °C for 1100 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The reaction was then diluted with 10 mL EtOAc, washed with an aqueous
- 10 solution of NH_4Cl , dried with anhydrous MgSO_4 , filtered and then the organic phase was evaporated *in vacuo*. The crude product was purified on silica column with 3:1 n-heptane:EtOAc as the mobile phase. Upon dissolving the crude product in the mobile phase, an insoluble precipitate was collected. On analysis this showed to be mainly pure product. All insoluble precipitate was dissolved in acetone, celite TM was added,
- 15 whereafter the acetone was evaporated. The celite was then applied to a silica column with 2:1 n-heptane:EtOAc as the mobile phase to give 34 mg (62%) of 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile as beige crystals. $M/Z = 258$.

Example 19

20




4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile

- 25 4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was coupled with (1-amino-cyclopentyl)-methanol (32 mg, 0.275 mmol), and DIPEA (36 mg, 0.275 mmol) in DMSO 0.7 mL using the same procedure as described in Example-8. This gave 23 mg (38%) of 4-(1-hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile as white powder. $M/Z = 284$.

30

5

Example 20




(S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile

10

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was coupled with (S)-2-amino-butan-1-ol (25 mg, 0.275 mmol), DIPEA (36 mg, 0.275 mmol), in 0.7 mL DMSO using the same procedure as described in Example-8. This gave 17 mg (31%) of (S)-4-(1-hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile as white crystals. M/Z

15 = 258.

Example 21




20 (R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.21 mmol), (R)-2-Amino-pentan-1-ol (32 mg, 0.27 mmol) and DIPEA (47 μ L, 0.27 mmol) was dissolved in DMSO (1 mL) and heated to 180 °C for 900 seconds in a microwave oven (Parameters: Fixed Holding time, High absorbance, pre-stirring 25 sec.). The crude product was diluted with CH_2Cl_2 and washed with an aqueous solution of NH_4Cl . The organic phase was separated, dried and evaporated in vacuo. The crude product was purified on a silica column with 3:1 n-heptane: EtOAc as the mobile phase. This gave 39 mg (68%) of (R)-4-(1-hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile. M/Z = 272.

25

5


Example 22

10

(S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.21 mmol) was coupled with (S)-2-Amino-pentan-1-ol (32 mg, 0.27 mmol), DIPEA (47 μ L, 0.27 mmol) in DMSO 1.0 mL, using the same procedure as described in Example-21. This gave 24 mg (42%) of (S)-4-(1-hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile. $M/Z = 272$

15

Example 23

20

[4-(R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile


(4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (100 mg, 0.492 mmol) was dissolved in DMSO (3.5 mL) and (R)-(-)-2-Amino-1-pentanol (66 mg, 0.634 mmol) and pyridine (52 μ L, 0.634 mmol) was added. The reaction was heated in microwave to 170 $^{\circ}$ C for 900 sec (Parameters: 30 seconds pre-stirring, holding time on, normal absorption). The mixture was diluted with EtOAc and washed with aqueous solution of NH_4Ac . The water phase was washed with EtOAc and the organic phases were pooled, dried with MgSO_4 , filtered

25

- 5 and evaporated *in vacuo*. The crude product was purified on a silica column with 5:1 *n*-heptane: EtOAc as the mobile phase. This gave 2.1 mg (1.5 %) of [4-(R)-1-hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. $M/Z = 286$

Example 24

10




[4-(S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile

- 15 (4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (100 mg, 0.492 mmol) was coupled with (S)-(+)-2-Amino-1-pentanol (66 mg, 0.634 mmol), Pyridine (52 μ L, 0.634 mmol), in DMSO (3.5 mL) using the same procedure as described in Example-23. This gave 2.2 mg (1.6 %) of [4-(S)-1-hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. $M/Z = 286$

20

Example 25




- 25 [4-(S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile

(4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (119 mg, 0.584 mmol) was coupled with L-Leucinol (89 mg, 0.759 mmol), Pyridine (62 μ L, 0.759 mmol), DMSO (3.2 mL) using

- 5 the same procedure as described in Example-23. This gave 2.6 mg (1.5 %) of [4-((S)-1-hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. $M/Z = 300$

Example 26



10

4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile


- 15 The 2-methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol (360 mg, 1.6 mmol) was dissolved in ethanol (26 ml) and $\text{Na}_2\text{S}_2\text{O}_4$ (2.23 g, 12.8 mmol) was added and the solution heated at 80°C overnight. The solvent was evaporated and the remaining solid was partitioned between 10% aqueous solution NaHCO_3 and EtOAc. The water phase (pH = neutral) was extracted with EtOAc (x 3), the collected organic phase washed with brine and dried (MgSO_4). The 2-(4-amino-3-methyl-phenylamino)-2-methyl-propan-1-ol was
- 20 used in the next step without further purification. (The amine oxidises on the TLC plate; brown spots upon standing.)

- 25 Sodium nitrite (NaNO_2) (190 mg, 2.75 mmol) in water (2.5 ml) was added to a solution of amine (500 mg, 2.5 mmol conc. HCl/ice (2.5 ml/2.5 g) during 5 min. followed by neutralisation by addition of solid CaCO_3 . KCN (391 mg, 6 mmol) and CuCN (269 mg, 3.0 mmol) in water (1 ml) was heated at 60°C (oil bath) and the cold, neutral diazonium salt solution was added drop wise over 15 min. Gas evolution was observed and the resulting suspension turned bright and strong orange. The reaction mixture was heated at 110°C for 30 min, cooled, diluted with water and EtOAc and filtered through celite. The
- 30 water phase was extracted with EtOAc and the collected organic phase washed with brine and dried (MgSO_4). The crude product (491 mg) was purified by flash chromatography

- 5 (Hexane; Hex/EtOAc; 7:3 → 1:1) giving the reduced compound 2-methyl-2-(3-hydroxy-phenylamino)-propan-1-ol (93 mg) and 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile (108 mg, 21%) as a pale yellow solid. M/Z = 204.

Example 27

10




6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile

- 15 2-Methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol (1.08 g, 4.8 mmol) was dissolved in 75% aqueous ethanol and $\text{Na}_2\text{S}_2\text{O}_4$ (3.9 g, 24 mmol) was added in portions. The reaction mixture was heated at 60°C for 30 min when TLC (10% MeOH in DCM) showed full conversion. The heat was turned off, the reaction mixture stirred overnight at ambient temperature and evaporated to dryness. To this residue was added NaHCO_3 (5% aq.) and EtOAc, the phases separated and the water phase (pH 7-8) extracted extensively
- 20 with EtOAc. (The product is very water-soluble and it is probably better to do a continuous extraction with EtOAc to get a higher yield). The collected organic phase was washed with brine before drying (MgSO_4). Upon standing, the colour of the organic solution turned from yellow to orange. Filtration and evaporation yielded 0.648 g (69%) of amine
- 25 as a red oil.

- NaNO_2 (0.25 g, 3.65 mmol) in water (3 ml) was added to a solution of amine 6 (0.648 g, 3.3 mmol) in ice/conc. HCl (3.5 g/3.5 ml) during 5 min. followed by neutralisation by addition of solid CaCO_3 . KCN (0.52 g, 7.96 mmol) and CuCN (0.36 g, 3.98 mmol) in
- 30 water (3 ml) was heated at 60°C (oil bath) and the cold, neutral diazoniumsalt solution was added drop wise over 15 min. Gas evolution was observed and the resulting

- 5 suspension turned bright and strong orange. The reaction mixture was heated at 110°C for 30 min, cooled, diluted with water and EtOAc and filtered through celite. The water phase was extracted with EtOAc and the collected organic phase was washed with brine and dried (MgSO₄). The crude product (0.248 g) was purified by flash chromatography (Hexane → Hex :EtOAc 3:7) yielding 34 mg of 2-methyl-2-(6-methyl-pyridin-2-ylamino)-propan-1-ol and 11 mg of 6-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile, M/Z = 205.

Example 28



15

4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile

- 20 The nitro compound 18 (0.20 g, 0.84 mmol) was dissolved in EtOH (20 ml), Na₂S₂O₄ (1.1 g, 6.71 mmol) was added and the reaction mixture heated at 80°C overnight. The cold reaction mixture was filtered through celite, washed well with EtOAc and the filtrate evaporated to dryness. The crude 2-(4-amino-2, 3-dimethyl-phenylamino)-2-methyl-propan-1-ol (0.292 g), pure by ¹H-NMR, was used as such in the next reactions.
- 25 The reaction was performed using the same procedure as described in Example-21 using 2-(4-amino-2,3-dimethyl-phenylamino)-2-methyl-propan-1-ol (0.175 g, 0.84 mmol) in conc. HCl/ice water (1 ml/5 ml), NaNO₂ (64 mg mg, 0.92 mmol) in water (1 ml), KCN (130 mg, 2 mmol) and CuCN (90 mg, 1 mmol) in water (1 ml). The crude product (341 mg) was purified by flash chromatography (Hexane; Hex 7/EtOAc 3) giving reduced
- 30 compound 2-(2,3-dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol and 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile. All the fractions containing impure nitrile were collected and crystallised from hexane/EtOAc to give 25

- 5 mg (13%) of pure 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile.
M/Z = 218.

Procedure for Library synthesis (Examples 29-86).


- 10 The following is the general procedure for library synthesis for the examples of 29-88.
The compounds are shown in table 2.

Smith-vials for the microwave oven were charged with 0.1 mmol either of the starting materials; 5-fluoro-2-nitro toluene, 5-fluoro-2-nitrobenzotrifluoride, 6-fluoro-2-methyl-3-

- 15 nitro-pyridine.










To each vial was added 0.5 ml DMSO, 20 μ L triethylamine (1.4 equivalents), and 1.4 equivalents of the diverse amino alcohols. The vials were run 1100s in 140°C in a microwave oven. After synthesis the products were analysed by LC-MS. The DMSO solutions were transferred to test tubes, and evaporated onto silica gel under reduced pressure. The silica gel from the tubes was placed on SPE SI columns, and a frit was placed on top. The products were purified with a gradient solution of heptane/EtOAc. The fractions were pooled and solvent was evaporated. Compounds which were more than 90% pure were tested in an *in vitro* assay which is described below. Purity was determined by analytic HPLC.

The scaffold used for the construction of the library is according to Formula II. The






















Formula II

Example	R ⁹	R ⁶	Z	Yield (%)	MS (-Q1)
29		CF ₃	CH	46	262.9
30		CF ₃	CH	55	290.8
31		CF ₃	CH	24	249.1
32		CF ₃	CH	62	276.7
33		CF ₃	CH	65	290.8
34		CF ₃	CH	23	290.8
35		CF ₃	CH	93	325.3
36		CF ₃	CH	78	341.2

Example	R9	R6	Z	Yield (%)	MS (-Q1)
37		CF ₃	CH	82	262.9
38		CF ₃	CH	95	305.2
39		CF ₃	CH	98	323.2
40		CF ₃	CH	98	290.8
41		CF ₃	CH	89	385
42		CF ₃	CH	92	290.8
43		CF ₃	CH	95	290.8
44		CF ₃	CH	100	378.1
45		CF ₃	CH	84	316

Example	R9	R6	Z	Yield (%)	MS (-Q1)
46		CF ₃	CH	90	262.9
47		CF ₃	CH	106	275.2
48		CF ₃	CH	75	304.3
49		CF ₃	CH	69	275.2
50		CF ₃	CH	76	370
51		CF ₃	CH	89	325.3
52		CH ₃	N	53	238.0
53		CH ₃	N	53	238.0
54		CH ₃	N	30	195.7

Example	R ₉	R ₆	Z	Yield (%)	MS (-Q1)
55		CH ₃	N	60	223.9
56		CH ₃	N	63	238.0
57		CH ₃	N	22	238.0
58		CH ₃	N	88	272.2
59		CH ₃	N	65	209.8
60		CH ₃	N	60	252.1
61		CH ₃	N	79	252.1
62		CH ₃	N	89	252.1
63		CH ₃	N	74	270.4

Example	R9	R6	Z	Yield (%)	MS (-Q1)
64		CH ₃	N	84	238.0
65		CH ₃	N	78	332.2
66		CH ₃	N	88	238.0
67		CH ₃	N	80	224.2
68		CH ₃	N	75	238.0
69		CH ₃	N	72	209.8
70		CH ₃	N	58	223.1
71		CH ₃	N	52	222.1
72		CH ₃	N	90	272.2
73		CH ₃	C	44.0	208.9

Example	R9	R6	Z	Yield (% MS (-Q1))	
74		CH ₃	CH	55.0	237.1
75		CH ₃	CH	66.0	195.1
76		CH ₃	CH	31.0	237.1
77		CH ₃	CH	30.0	223
78		CH ₃	CH	32.0	237.1
79		CH ₃	CH	27	271.3
80		CH ₃	CH	25	250.9
81		CH ₃	CH	27	269.2
82		CH ₃	CH	24	237.1
83		CH ₃	CH	24	237.1
84		CH ₃	CH	24	237.1




Example	R9	R6	Z	Yield (%)	MS (-Q1)
85		CH ₃	CH	25	250
86		CH ₃	CH	33	316

Table 2


Example 87

15 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid

A suspension of 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile (70 mg, 0.34 mmol) and NaOH (0.14 g, 3.4 mmol) in water/MeOH (5 ml/8 ml) was refluxed for 4 days. The reaction mixture was diluted with water, pH adjusted to approx. 3 with 50% aq. HCl. The precipitated solid was filtered off and collected, the water phase was extracted with EtOAc (x 3), washed with brine and dried (MgSO₄). The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 39 mg (51%)

- 5 of the 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid as a brownish foam. M/Z 223.

Example-88




10

(6-Methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester

- 6-chloro-3-nitro-2-picoline (600 mg, 3.5 mmol) was coupled with glycine methyl ester hydrochloride (880 mg, 7 mmol), triethylamine (1.5 ml, 10.5 mmol) in DMSO 3 mL at 15 140 °C for 30 min in microwave (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The crude mixture was treated with a saturated aqueous solution of NH₄Cl. The aqueous solution of was extracted with EtOAc, washed with water and brine. The crude product was purified on a silica column with CH₂Cl₂-MeOH as mobile phase. This gave 39 mg (51%) of 580 mg (74%) of (6-methyl-5-nitro-2-pyridin-2-ylamino)- 20 butionic methyl ester as a yellow solid. M/Z = 225.

Example-89



- 25 2-Methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol

2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butionic methyl ester (30 mg, 0.13 mmol) was dissolved in THF (3 mL) and added to a nitrogen-purged flask containing methyl magnesium chloride (MeMgCl) (0.08 mL, 0.027 mmol) at 0 °C. The reaction mixture was

- 5 allowed to reach room temperature and then refluxed for 5 h. The reaction was quenched by adding saturated NH_4Cl . The reaction mixture was extracted with EtOAc and washed with H_2O and brine. The crude product was purified by HPLC. This gave 1.5 mg (5%) of 2-methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol as yellow oil. $M/Z = 225$.

10

Example-90

AR Competition Binding Assay

- Recombinant human androgen receptor (hAR) was extracted from Sf9 insect cells with
 15 buffer containing 1 mM EDTA, 20 mM K_2HPO_4 , 8.7% glycerol, 20 mM Na_2MoO_4 and 12 mM MTG at 5×10^7 cells/ml. The cell debris was removed by centrifugation and the supernatant aliquoted and stored at -70°C .
 An aliquot of AR extract was thawed on ice prior to use and diluted to approximately 0.2
 nM (1 to 30 dilution) in buffer (100 mM $\text{K}_n\text{H}_m\text{PO}_4$ pH 7.0, 1 mM EDTA, 8.7% glycerol,
 20 20 mM Na_2MoO_4 and 1 mM DTT). The test ligands were diluted in DMSO as a dilution series of 10 concentrations in duplicate, with 1:5 dilution between each concentration.
 Tritiated mibolerone (^3H -Mib) was used as tracer compound and diluted to 1.6 nM in 1
 mM EDTA, 20 mM Na_2MoO_4 , 8.7% glycerol and 1 mM DTT. To a 96-well
 polypropylene-plate 110 μl /well of 1.6 nM ^3H -Mib, 10 μl /well test substance and 110
 25 μl /well diluted AR was added. The plates were covered and incubated at $+4^\circ\text{C}$ over night.
 The plates were harvested on filters to separate bound ligand from unbound ligand with a
 Tomtec Harvester. A prewet buffer containing 20 mM $\text{K}_n(\text{PO}_4)$ pH 7.6, 1 mM EDTA, v/v
 0.5% polyethylenimine was used to equilibrate the filter before filtering the samples and
 washing the filters with 20 mM $\text{K}_n(\text{PO}_4)$ pH 7.6, 1 mM EDTA 8 times. The filters were
 30 allowed to dry for 1 hour at $+65^\circ\text{C}$. A scintillating wax was melted upon the filter and the
 radioactivity retained on the filter was measured in a Wallac Microbeta scintillation
 counter.

- 5 The affinity to AR was evaluated by a non-linear four-parameter logistic model: $b = (b_{\max} - b_{\min}) / (1 + (IC_{50}/I)^S) + b_{\min}$, where b_{\max} = total concentration of binding sites, b_{\min} = non-specific binding, I = added concentration of binding inhibitor, IC_{50} = concentration of binding inhibitor at half-maximal binding and S = slope factor.
- 10 Table: Antagonist and partial antagonist and binding activity of androgen receptor modulator compounds.

AR Transactivation Assays

- 15 The agonist and antagonist properties of compounds were determined using a cell-based system expressing stably integrated androgen receptor and an androgen responsive reporter gene. CV-1 cells (kidney fibroblasts) stably expressing CMV-hAR and alkaline phosphatase (ALP) driven by an MMTV promoter containing an androgen response element were cultured in Dulbecco's Modified Eagle Medium (DMEM), low glucose supplemented with 10% fetal bovine serum, 1% L-glutamine, and 0.7% Hygromycin B.
- 20 The stably integrated cells (ARAF) were trypsinized and resuspended in Opti-MEM 1 supplemented with 2% fetal bovine serum, 1% L- Glutamine, 50 µg/ml Gentamicine and 1% Pen/Strep. The cells were counted in a Birch chamber and diluted to a concentration of 100 000 cells /ml. The cells were then seeded out in 384 plates, 5000cells/well in 50µl seeding media and incubated overnight in 37 C, 5% CO₂.
- 25 The next day, the seeding medium was removed from the cells and 20 µl induction media (Opti-MEM 1 supplemented with 1% L- Glutamine, 50 µg/ml Gentamicine and 1% Pen/Strep) +/- 0.1 nM Mibolerone was added to the wells. 10µl of test compound diluted in induction media was then added to the wells. The cells were incubated 48 hr in 37 C, 5% CO₂.
- 30 After 48 hr 5µl of cell medium was added to white 384 plates with 100µl of ALP substrate buffer. The plates were incubated in 37 C for 20 minutes followed by incubation at room temperature for 10 minutes before each well was read in a µBETA machine. Agonist activity was calculated from the alkaline phosphatase activity induced

- 5 in the absence of Mibolerone and compared to standard activation curve generated by Mibolerone alone. Antagonist activity was calculated from the decrease in ALP activity in the presence of 0.1 nM Mibolerone. EC50 and IC50 values were calculated by using a non-linear four-parameter fit as described above.
- 10 Other assays to determine androgen receptor mediated activity of the test compounds include modulation of endogenous AR mediated transcription in cell culture systems; modulation of androgen responsive tissue effects in rodents; identification of receptor surface conformation changes; and binding specificity to AR versus other nuclear receptors.

15

	AR_LT IC50 (nM)	ARAF EC50 (nM)	ARAF %AGONIST	ARAF IC50 (nM)	ARAF %ANTAGONIST
Example-1	22.77	26.8	51.7	2.1	33.1
Example-5	38.06	81.7	29.3	7.2	61
Example-8	241.44	374.2	10.6	22.3	82.5
Example-19	130.38			22.4	95.6
Example-30	113.45	1069.9	7.3	61.8	90.9
Example-41	7.5			26.7	98.2
Example-42	127			23.4	99.2
Example-60	14			3.2	90.3
Example-69	492.4	nt	nt	nt	nt
Example-77	44.5	nt	nt	nt	nt
Example-86	74.1			71.1	100


nt = not tested

20

CLAIMS

5

1. Use as a medicament of a compound according to Formula I



Formula I

10

in which;

- 15 R_1 and R_2 are the same or different and independently selected from the group consisting of; hydrogen, halogen, C_1 - C_{10} alkyl, C_3 - C_{10} cycloalkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkoxy, C_1 - C_{10} alkenoxy, C_1 - C_{10} alkynoxy, C_1 - C_{10} alkylthio, C_1 - C_{10} alkenylthio, C_1 - C_{10} alkynylthio, C_6 - C_{10} arylthio, C_1 - C_{10} alkylsulphone, C_1 - C_{10} alkenylsulphone, C_1 - C_{10} alkynylsulphone, C_6 - C_{10} arylsulphone, C_1 - C_{10} alkylsulphoxide, C_1 - C_{10} alkenylsulphoxide, C_6 - C_{10} arylsulphoxide, C_6 - C_{10} aryl, or C_5 - C_{20} heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be
20 the same or different;

- 25 R_3 and R_4 are the same or different and independently selected from the group consisting of; hydrogen, halogen, keto, C_1 - C_{20} alkyl, C_3 - C_7 cycloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 alkenoxy, C_1 - C_4 alkynoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkenylthio, C_1 - C_4 alkynylthio, C_1 - C_{10} alkylsulphone, C_1 - C_{10} alkenylsulphone, C_1 - C_{10} alkynylsulphone, C_6 - C_{10} arylsulphone, C_1 - C_{10} alkylsulphoxide, C_1 - C_{10} alkenylsulphoxide, C_6 - C_{10} arylsulphoxide, C_6 - C_{15} aryl, or C_5 - C_{20} heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be
the same or different;

5

R_5 is chosen from the group consisting of: nitro, cyano, acetic acid, halogen, sulphonic acid, aldehyde, carboxylic acid or ester, phosphonic acid, or ester;

10

R_6 is from the group consisting of: C_1 - C_5 alkyl, halogen, CN, CO_2H , CHF_2 , CH_2F , or CF_3 ;

R_8 is chosen from the group consisting of: H, C_1 - C_5 alkyl, halogen, CHF_2 , CH_2F or CF_3 ;

X is chosen from the group consisting of: $-NH-$, $-O-$, $-S-$, $-SO-$, $-SO_2$, $-Se-$, $-Te-$ or $-S-S-$

15

Y is chosen from the group consisting of: OH, NH_2 , unbranched, branched or cyclic C_1 - C_5 alkyl, unbranched, branched or cyclic $-NH(C_1-C_8)$; unbranched, branched or cyclic $N(C_1-C_8)_2$, $-NH(C_6 \text{ aryl})$, $-N(C_6 \text{ aryl})_2$, $-NH(C_1-C_5 \text{ heteroaryl})$, and $-N(C_1-C_5 \text{ heteroaryl})_2$ or a bioisosteric equivalent;

20

Z is chosen from the group consisting of: CR_7 , N, or O;

R_7 is H, or C_1 - C_5 alkyl;

25 R^a represents a member selected from the group consisting of: hydrogen, halogen, $-CN$, CO_2H , CHO , NO_2 , $-NH_2$, $-NH(C_1-C_4)$; $N(C_1-C_4)_2$, $-NH(C_6 \text{ aryl})$, $-N(C_6 \text{ aryl})_2$, $-NH(C_1-C_5 \text{ heteroaryl})$, and $-N(C_1-C_5 \text{ heteroaryl})_2$ or a bioisosteric equivalent; or a pharmaceutically acceptable salt thereof;

30

2. Use according to claim 1, wherein R_1 or/and R_2 are H, (S)-methyl, methyl, (R)-ethyl, (S)-ethyl, ethyl, (R)-propyl, (S)-propyl, propyl, (S)-butyl, (S)-1-methyl-propyl, (S)-2-methyl-propyl, (R)-isopropyl, (S)-isopropyl, isopropyl, cyclopentyl, $-(CH_2)_2SMe$, (R)- CH_2SCH_2Ph , (S)-benzyl, 4-chloro-benzyl, (S)-3-methyl-1-H-indole or (S)-phenyl;

5

3. Use according to either of the preceding claims wherein R_3 is chosen from the group consisting of; hydrogen, methyl, ethyl, phenyl, 3-hydroxy phenyl, 4-hydroxy phenyl, or keto.

10 4. Use according to any of the preceding claims wherein R_4 is H, methyl, or keto.

5. Use according to any of the preceding claims wherein R_5 is NO_2 , CN, CH_2CN or CO_2H ;

15 6. Use according to any of the preceding claims wherein R_6 is Me, or CF_3 ;

7. Use according to any of the preceding claims wherein R_7 is H or Me;

8. Use according to any of the preceding claims wherein R_8 is H or methyl;

20

9. Use according to any of the preceding claims wherein X is NH ;

10. Use according to any of the preceding claims wherein Y is H, -OH, -OMe, -N $(\text{CH}_2\text{CH}_3)_2$, piperidine, or 4-nitro-2-ylamino;

25

11. Use according to any of the preceding claims wherein Z is CR_7 or N;


12. Use according to any of the preceding claims wherein the compound is chosen from the group consisting of;

30

2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol;
[1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol;
(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol;










- 5 (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol;
2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol;
[1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol;
(S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol;
2-Methyl-2-(6-methyl-5-nitro-pyridine-2-ylamino)-propan-1-ol;
- 10 [1-(6-Methyl-5-nitro-pyridine-2-ylamino)-cyclopentyl]-methanol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-2-phenyl-ethanol;
(S)-2-(6-Methyl-5-nitro-pyridine-2-ylamino)-3-phenyl-propan-1-ol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol;
(DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
- 15 (S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol;
(S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile;
- 20 4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
(S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
[4-((S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
- 25 [4-((R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile;
6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile;
- 30 and compounds having the formula:










5













in which R_9 , R_6 and Z are as defined in the following table:

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH

R9	R6	Z		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	CH

[illegible]



- 5 15. Use of a compound as defined in Formula I of any preceding claim for manufacturing a medicament to be used in the treatment of a disease caused by a disturbance in the activity of the androgen receptor.
16. Use according to claim 15 wherein the disease is caused by an increase in androgen
- 10 receptor activity.
17. Use according to claim 15 or 16 wherein the disease is chosen from the group consisting of, prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic
- 15 hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.

20

5

ABSTRACT

The present invention relates to novel compounds which are active as antagonists or
10 partial antagonists of the androgen receptor, and to the use of such compounds in the field
of medicine. Furthermore, the compounds can be used for preparing medicaments for
treating diseases related to disturbances in the activity of the androgen receptor.

15